



Five Things Physicians and Patients Should Question

1

Don't perform routine pre-operative hemostatic testing (PT, aPTT) in an otherwise healthy child with no prior personal or family history of bleeding.

Preoperative hemostatic screening in healthy pediatric patients with no personal or family history of excessive bleeding does not effectively identify those who will have unexpected surgical bleeding. Artifacts or disorders that do not affect bleeding risk may be identified, such as factor XII deficiency or an infection-associated, transient lupus anticoagulant. Hemostatic testing adds cost and may introduce additional stress, either due to blood sampling or if a child has "abnormal" results.

2

Don't transfuse platelets in an asymptomatic (i.e., non-bleeding) pediatric patient (e.g. aplastic anemia, leukemia, etc.), with a platelet count > 10,000/mcL unless other signs and/or symptoms for bleeding are present, or if the patient is to undergo an invasive procedure.

In asymptomatic (i.e., non-bleeding) pediatric patients with a platelet count > 10,000/mcL, transfusion is not clinically indicated unless signs, symptoms, or increased risk factors of bleeding are present. This practice is consistent with recommendations from the clinical guidelines of multiple associations (National Institute for Health and Care Excellence, British Society for Haematology, American Society of Clinical Oncology, and American Society of Hematology). The risk of spontaneous bleeding is low at platelet counts > 10,000/mcL. Unnecessary transfusions put patients at risk for transfusion reactions, alloimmunization, blood borne infections, and refractoriness to future platelet transfusions. This recommendation does not apply in anticipation of an invasive procedure.

3

Don't order thrombophilia testing on children with venous access (i.e., peripheral or central) associated thrombosis in the absence of a positive family history.

Testing for inherited forms of thrombophilia does not influence the initial management of a first episode of provoked venous thrombosis and should not be performed routinely. The results of such testing have not been shown to either predict recurrence of venous thrombosis or inform the intensity or duration of anticoagulant therapy. Thrombophilia testing has substantial financial cost, and a positive result has the potential for misinterpretation of risk assessment leading to undue psychological distress or impact on childbearing plans, as well as possible life insurance discrimination for affected patients.

4

Don't transfuse packed red blood cells (pRBC) for iron deficiency anemia in asymptomatic pediatric patients when there is no evidence of hemodynamic instability or active bleeding.

In pediatric patients with asymptomatic, iron deficiency anemia, do not transfuse packed red blood cells (pRBC) in the absence of hemodynamic instability or active bleeding. Unnecessary pRBC transfusions put patients at risk for complications, such as transfusion reactions, blood borne infections and volume overload. The judicious use of pRBCs transfusions would also be associated with cost savings for healthcare systems.

5

Don't routinely administer granulocyte colony stimulating factor (G-CSF) for empiric treatment of pediatric patients with asymptomatic autoimmune neutropenia in the absence of recurrent or severe bacterial and/or fungal infections.

In pediatric patients with asymptomatic autoimmune neutropenia, there is insufficient evidence to support the routine use of granulocyte colony stimulating factor (G-CSF) as a prophylaxis strategy to improve health outcomes. The use of G-CSF in this population should be guided by clinical evaluation. The unnecessary routine use of G-CSF could lead to intolerable side effects, such as bone pain, as well as avoidable healthcare cost.

How This List Was Created

The American Society of Hematology (ASH) and the American Society of Pediatric Hematology/Oncology (ASPHO) formed a task force to solicit, evaluate, and select list items for a pediatric focused Choosing Wisely list. The panel was composed of 13 members – two co-chairs (representing ASH and ASPHO), five members selected by each organization, and one member serving as an advisor on Choosing Wisely methodology. Suggestions were solicited from the membership of both societies. Formal systematic reviews of the evidence were completed for eight semi-finalist items. Final item selections were made by the ASH-ASPHO CW task force with reference to the following six guiding principles: avoiding harm to patients, producing evidence-based recommendations, considering both the cost and frequency of tests and treatments, making recommendations in the clinical purview of the hematologist, and considering the potential impact of recommendations. Harm avoidance was established as the campaign's preeminent guiding principle.

Sources

- Alzahrani A, Othman N, Bin-Ali T, Elfaraidi H, Al Mussaed E, Alabbas F, Sedick Q, Albatniji F, Alshahrani Z, Asiri M, Alsuhaibani O, Elyamany G. Routine Preoperative Coagulation Tests in Children Undergoing Elective Surgery or Invasive Procedures: Are They Still Necessary? *Clinical Medicine Insights: Blood Disorders*. 2019 Jan;12:1-4.
Bidlemaier C, Olivieri M, Hütker S, Dietl S, Kurnik K. Perioperative management of hemostasis in children and adolescents. *Blood Cells, Molecules, and Diseases*. 2017 Jan;67:91-95.
Chee YL, Crawford JC, Watson HG, Greaves M. Guidelines on the assessment of bleeding risk prior to surgery or invasive procedures British Committee for Standards in Haematology. *British Journal of Haematology*. 2008 Feb;140:496-504.
Cooper JD, Smith KJ, Ritchey AK. A cost-effectiveness analysis of coagulation testing prior to tonsillectomy and adenoidectomy in children. *Pediatric Blood & Cancer*. 2010 Dec;55(6):1153-1159.
Cosmi B, Alatri A, Cattaneo M, Gesele P, Marietta M, Rodeghiero F, Tripodi A, Ansaloni L, Fusari M, Taddei S. Assessment of the risk of bleeding in patients undergoing surgery or invasive procedures: Guidelines of the Italian Society for Haemostasis and Thrombosis (SIST). *Thrombosis Research*. 2009 Nov;124(5):e6-e12.
Lescanne E, Chiron B, Constant I, Couloigner V, Fauroux B, Hassani Y, Jouffroy L, Lesage V, Mondain M, Nowak C, Orliaguet G, Viot A. Pediatric tonsillectomy: Clinical practice guidelines. *European Annals of Otorhinolaryngology, Head and Neck Diseases*. 2012 Oct;129(5):264-271.
van Veen JJ, Spahn DR, Makris M. Routine preoperative coagulation tests: an outdated practice? *British Journal of Anaesthesia*. 2011 Jan;106(1):1-3.
- National Institute for Health and Care Excellence. Blood transfusion NICE Guideline 24. 2015 Nov; Retrieved from www.nice.org.uk/guidance/ng24.
Neunert C, Lim W, Crowther M, Cohen A, Solberg L Jr, Crowther MA. The American Society of Hematology 2011 evidence-based practice guideline for immune thrombocytopenia. *Blood*. 2011 Apr;117(16):4190-4207.
New HV, Berryman J, Bolton-Maggs PH, Cantwell C, Chalmers EA, Davies T, Gottstein R, Kelleher A, Kumar S, Morley SL, Stanworth SJ. Guidelines on transfusion for fetuses, neonates and older children. *British Journal of Haematology*. 2016 Nov;175(5):784-828.
Schiffer CA, Bohlke K, Delaney M, Hume H, Magdalinski AJ, McCullough JJ, Omel JL, Rainey JM, Rebutta P, Rowley SD, Troner MB, Anderson KC. Platelet Transfusion for Patients With Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update. *Journal of Clinical Oncology*. 2018 Jan;36(3):283-299.
- Baglin T, Gray E, Greaves M, Hunt BJ, Keeling D, Machin S, Mackie I, Makris M, Nokes T, Perry D, Tait RC, Walker I, Watson H. Clinical guidelines for testing for heritable thrombophilia. *British Journal of Haematology*. 2010 Apr;149(2):209-220.
Chalmers E, Ganesh V, Liesner R, Maroo S, Nokes T, Saunders D, Williams M. Guideline on the investigation, management and prevention of venous thrombosis in children. *British Journal of Haematology*. 2011 Jul;154(2):196-207.
National Institute for Health and Care Excellence. Venous thromboembolic diseases: diagnosis, management and thrombophilia testing. 2015 Nov; Retrieved from www.nice.org.uk/guidance/cg144.
Scottish Intercollegiate Guidelines Network. Prevention and management of venous thromboembolism. 2014 Oct; Retrieved from www.sign.ac.uk/sign-122-prevention-and-management-of-venous-thromboembolism.html.
- Alberta Medical Association. Iron Deficiency Anemia (IDA). 2018 Mar; Retrieved from actt.albertadoctors.org/CPGs/Lists/CPGDocumentList/IDA-CPG.pdf.
British Columbia Guidelines and Protocols Advisory Committee. Iron Deficiency – Diagnosis and Management. 2019 April; Retrieved from www2.gov.bc.ca/gov/content/health/practitioner-professional-resources/bc-guidelines/iron-deficiency.
- Fioredda F, Calvillo M, Bonanomi S, Coliva T, Tucci F, Farruggia P, Pillon M, Martire B, Ghilardi R, Ramenghi U, Renga D, Menna G, Pusiol A, Barone A, Gambineri E, Palazzi G, Casazza G, Lanciotti M, Dufour C. Congenital and acquired neutropenias consensus guidelines on therapy and follow-up in childhood from the Neutropenia Committee of the Marrow Failure Syndrome Group of the AIEOP (Associazione Italiana Emato-Oncologia Pediatrica). *American Journal of Hematology*. 2012 Feb;87(2):238-243.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.

To learn more about the ABIM Foundation, visit www.abimfoundation.org.



About the American Society of Hematology

The American Society of Hematology (ASH) is the world's largest professional society of hematologists, serving more than 17,000 clinicians and scientists from around the world who are dedicated to furthering the understanding, diagnosis, treatment and prevention of disorders affecting the blood.

For more than 50 years, the Society has led the development of hematology as a discipline by promoting research, patient care, education, training and advocacy in hematology. By providing a forum for clinicians and scientists to share the latest discoveries in the field, ASH is helping to improve care and possibly lead to cures for diseases that affect millions of people, including leukemia, lymphoma, myeloma, anemias and various bleeding and clotting disorders.

For more information, visit www.hematology.org.

About the American Society of Pediatric Hematology/Oncology

The American Society of Pediatric Hematology/Oncology (ASPHO) is the professional medical society of pediatric hematology/oncology subspecialists and other healthcare professionals dedicated to promoting the optimal care of children, adolescents and young adults with blood disorders and cancer. Founded in 1981, ASPHO serves over 2000 members including physicians, researchers, advanced practice providers, and fellows in training by sponsoring educational and professional development programs, promoting discovery, advocating on behalf of its patient base and members, advancing professional practice and supporting partnerships to further its goals.

For more information, visit www.aspho.org.



For more information or to see other lists of Things Providers and Patients Should Question, visit www.choosingwisely.org.